

## Critically Appraised Topic (CAT)

<b>Project Team:</b>
8A2
<b>Project Team Participants:</b>
Ardit Haxia, Matthew Boeker, Megan Hunjadi, Claudia VanOpdorp
<b>Clinical Question:</b>
What is the progression and maintenance of salivary gland disease and its effect on dental management
<b>PICO Format:</b>
<b>P:</b>
Patients with submandibular sialolithiasis
<b>I:</b>
Less invasive techniques such as transoral stone removal, edoscopy or lithotripsy
<b>C:</b>
Submandibular gland resection
<b>O:</b>
Better outcome for the patient
<b>PICO Formatted Question:</b>
In patients with submandibular sialolithiasis, do less invasive techniques such as transoral stone removal, edoscopy or lithotripsy versus submandibular gland resection result in better outcomes for the patients?
<b>Clinical Bottom Line:</b>
<a href="#">Click here to enter text.</a>
<b>Date(s) of Search:</b>
11/02/20
<b>Database(s) Used:</b>
PubMed
<b>Search Strategy/Keywords:</b>
Sialolithiasis, submandibular gland resection, edoscopy
<b>MESH terms used:</b>
Endoscopy, Submandibular gland disease, Treatment outcome, Human
<b>Article(s) Cited:</b>
1. Zenk J, Koch M, Klintworth N, König B, Konz K, Gillespie MB, Iro H. Sialendoscopy in the diagnosis and treatment of sialolithiasis: a study on more than 1000 patients. Otolaryngol Head Neck Surg. 2012 Nov;147(5):858-63. doi: 10.1177/0194599812452837. Epub 2012 Jun 29. PMID: 22753615.

2. Jadu, Fatima M, and Ahmed M Jan. "A meta-analysis of the efficacy and safety of managing parotid and submandibular sialoliths using sialendoscopy assisted surgery." *Saudi medical journal* vol. 35,10 (2014): 1188-94.

3. Xiao JQ, Sun HJ, Qiao QH, Bao X, Wu CB, Zhou Q. Advantages of submandibular gland preservation surgery over submandibular gland resection for proximal submandibular stones. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2018 May;125(5):e113-e117. doi: 10.1016/j.oooo.2017.12.009. Epub 2017 Dec 29. PMID: 29530607.

4. Eun, Y.G., Chung, D.H. and Kwon, K.H. (2010), Advantages of intraoral removal over submandibular gland resection for proximal submandibular stones. *The Laryngoscope*, 120: 2189-2192. doi:[10.1002/lary.21120](https://doi.org/10.1002/lary.21120)

**Study Design(s):**

- 1: Case Series (5)
- 2: Meta Analysis (1a)
- 3: Randomized Control Trials (1b)
- 4: Randomized Control Trials (1b)

**Reason for Article Selection:**

- 1. Gave percentages of success rates for different techniques and combinations of the techniques.
- 2. This was a meta-analysis that looked at some of the complications of total gland resection and also at the benefits and safety of conservative gland surgery.
- 3. This was a randomized control trial that directly compared sialendoscopy assisted conservative removal and traditional SMG resection for the removal of submandibular stones.
- 4.

**Article(s) Synopsis:**

1. This article looked at 1154 patients with either submandibular or paratoid salivary stones from 2003 to 2008. It examined the long term success rates of different gland saving techniques for stone removal, including transoral stone removal, extracorporeal shockwave lithotripsy, and sialendoscopy. Total gland removal was also looked at in the study if there was not long term success from the above mentioned techniques. In submandibular sialolithiasis, transoral stone removal was the most used option to remove the stone, with only 4% of cases needing submandibular gland removal. Extracorporeal shockwave lithotripsy was the most common technique used for paratoid stone removal, although edoscopy and inscisional technique were also used at least 20% of the time. The success rate was above 75% for all techniques with only 4% of patients needing paratidectomy. Endoscopy was successful on submandibular stones with a mean diameter of 4.9mm, endoscopy with transoral stone removal or ESWL was successful for submandibular stones with a mean diameter of 7.6mm and transoral stone removal was successful in submandibular stones that were a mean diameter of 9.1mm. The conclusions reached in this study found that transoral stone removal was the treatment of choice for

submandibular stones if they could not be mobilized within a few minutes via edoscopy. This study also said the ideal sites for transoral submandibular stone removal are the distal duct or the hilar region of the gland.

2. This study was a meta-analysis that looked at articles from 2003 to 2014 dealing with the safety and efficacy of different techniques of salivary stone removal. The article goes on to say that stones larger than 7mm should not be removed conservatively. Round stones are also more congenial to conservative removal than irregular stones are. Asymptomatic stones should be managed conservatively and stones located in the primary or distal portion of the duct have a better outcome with conservative removal than do those stones located in the hilum or gland substance. Salivary function normally remains ok, but if there is too much distension in the gland itself this can lead to salivary pooling which then necessitates gland removal. This study then goes on to describe in detail lithotripsy, sialendoscopy, gland preserving surgery and sialadenectomy. For lithotripsy, the authors say that it is best used on immobile stones or stones that are over 7mm in diameter. For sialendoscopy, they recommended stones that were smaller than 7mm unless you use it in conjunction with other techniques such as gland preserving surgery or sialendoscopy or lithotripsy. Pain and swelling are the 2 main side effects and the only contraindication is acute sialadenitis. This study again listed a 92% success rate for submandibular transoral stone removal, with pain, transient lingual nerve paresthesia, hemorrhage and ranula formation. Removal of the gland is only used if more conservative methods fail or if recurring stones appear. Complications include severing of the facial nerve, Frey syndrome, hematoma, seroma and facial asymmetry from a transcervical entry point. It can also be done intraorally, and this can cause damage to the lingual and hypoglossal nerve. All of the stones were at least 4mm large and most were in difficult locations to reach.

3. This study was a randomized case control study that looked at 40 patients with submandibular stones. The groups were divided into two groups of 20, with one group having sialendoscopy assisted surgery and the other having traditional SMG resection. For the sialendoscopy assisted surgery, the incision was extra oral because of the placement of the stone, but still did not lead to complete gland resection. The results stated that in the conservative group, after a 6 month follow up, there was no damage to the lingual nerve, swelling only lasted 1-2 days, and the facial morphology remained intact. The average size of the stone removed was 6.3 mm. In the resection group, swelling lasted for 3-5 days, all patients had some facial morphology change, certain patients reported dry mouth and 1 patient had lingual nerve damage. Pain scores were significantly higher post op for the resection group as well. The study recognized that stones should be removed via intraoral incisions if possible, but due to the location of the stone this is not always possible. In these cases, extra oral incisions with sialendoscopy is a very good choice of treatment.

4

Levels of Evidence: (For Therapy/Prevention, Etiology/Harm)

See <http://www.cebm.net/index.aspx?o=1025>

☐ **1a** – Clinical Practice Guideline, Meta-Analysis, Systematic Review of Randomized Control Trials (RCTs)

☐ **1b** – Individual RCT

☐ **2a** – Systematic Review of Cohort Studies

☐ **2b** – Individual Cohort Study

☐ **3** – Cross-sectional Studies, Ecologic Studies, “Outcomes” Research

☐ **4a** – Systematic Review of Case Control Studies

☐ **4b** – Individual Case Control Study

☐ **5** – Case Series, Case Reports

☐ **6** – Expert Opinion without explicit critical appraisal, Narrative Review

☐ **7** – Animal Research

☐ **8** – In Vitro Research

**Strength of Recommendation Taxonomy (SORT) For Guidelines and Systematic Reviews**

See article **J Evid Base Dent Pract 2007;147-150**

☐ **A** – Consistent, good quality patient oriented evidence

☐ **B** – Inconsistent or limited quality patient oriented evidence

☐ **C** – Consensus, disease oriented evidence, usual practice, expert opinion, or case series for studies of diagnosis, treatment, prevention, or screening

**Conclusion(s):**

[Click here to enter text.](#)